

## Abandoned farmlands lead to more forest fires in mainland Portugal

June 18th, 2017



June 18 2017. Large forest yres strike Portugal, northeast of Lisbon. Many people have been killed. This article has been published before on the ClimateChangePost. A lot of research has been done over the last years on the increasing risk of forest yres in southern Europe, including the impact of climate change. Read our articles on the Forest Fires page, f.i. our longread overview on 'Wildyres and climate change, a connection that's hard to deny'.

Before the 1980s ftres in mainland Portugal had never reached 10,000 ha of burned area in a single occurrence, according to available data on forest ftres. In the last decade two forest ftres occurred that covered an area of over 20,000 ha. The number of large forest ftres in Portugal is increasing, and the extent of these ftres is increasing as well. According to scientists a new era of forest ftres has begun in Portugal, and it started somewhere around 1987.

Climate change can be an important driver of greater ftre activity, but it's not the only one. In fact, in southern Europe climate change seems to be a far less important driver than socio-economic factors and changes in land use. Important socio-economic factors are depopulation and ageing of rural populace. Important changes in land use are agricultural abandonment, expansion of highly sammable tree species, and renewed growth of woody vegetation in cleared landscapes. Also, the success in suppressing small and medium size ftres is producing landscapes with higher sammability and more fuel loads that contribute to larger and more severe forest ftres.

Already degraded areas become more likely to be affected by ftres. These areas are characterized by herbaceous plants and shrubs, which are extremely susceptible to ftre because of their low moisture content and sammable fuel load. This is a vicious cycle. The rural exodus led to an increase in abandoned farmlands and growth of woody vegetation. This soil degradation as a result leads to more and larger ftres that consequently further degrade the soil. Many of these rural areas became liable to the occurrence of devastating ftres as a result of the high quantities of biomass accumulated over the years, which could fuel catastrophic ftres during the summer months. The proportion of uncultivated land, due to the population exodus from rural areas to coastal cities and other European countries since the 1950s, was the most important factor affecting burnt areas in Portugal.

In general, very large forest ftres represent a small fraction of the total number of forest ftres but are responsible for the overwhelming majority of the total burned area. Forest ftres in Portugal over 5000 ha in the period 1981 - 2010, for instance, only represent 1% of the total number of forest ftres with burnt areas over 100 ha, but were responsible for 15% of the total area burned. The majority of forest ftres were located in the northern region of Portugal, and the Faro district in the south.

Source: Ferreira-Leite et al., 2016. Natural Hazards 84: 1035-1053

Photo: Steve McCaig (www.stckr.com)